Absolute Convergence Test (ACT)

Given a series
$$\sum_{n=1}^{\infty} a_n$$
, if the Absolute Series $\sum_{n=1}^{\infty} |a_n|$ Converges, then the Original Series $\sum_{n=1}^{\infty} a_n$ Converges.

USED: When the Absolute Series is easier to analyze.

USED: To avoid analyzing negative signs, or maybe the Alternating Series Test.

BENEFITS: Handling series with positive terms is usually easier. Plus, you have more convergence tests available for positive termed series; think IT, CT, LCT.

WARNING: If you analyze the Absolute Series, and it is NOT convergent, then this convergence test is Inconclusive and not an option for use. It does not immediately follow up with a conclusion about Conditional Convergence without more work. All you know if the the A.S. diverges is that your original series is NOT Absolutely Convergent (A.C.)

NOTE: If the Original Series is already equal to the Absolute series, then Absolute Convergence is the same as Convergence.

NOTE: If the Original Series is not the same as the Absolute Series, then this ACT says that **Absolute Convergence implies Convergence**.

APPROACH:

- Given the original series, find the Absolute Series by ignoring any negative signs.
- Analyze the Absolute Series completely as before. Run all the details of *p*-Series Test, IT, CT, or LCT. Justify everything. Make a clear conclusion about the convergence of the Absolute Series.
- If the A.S. Converges, then you are done, and the O.S. Converges by ACT.
- If the A.S. diverges, you have **no** conclusion about the O.S. You are stuck running a Convergence test (likely AST) on the O.S.

EXAMPLES: Determine and state whether each of the following series **Converges** or **Diverges**. Name any Convergence test(s) that you use, and justify all of your work.

1.
$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n^5} \to \text{Check the Absolute Series } \sum_{n=1}^{\infty} \frac{1}{n^5}$$
 which is a Convergent *p*-Series with $p = 5 > 1$. Then the Original Series $\sum_{n=1}^{\infty} (-1)^n \frac{1}{n^5}$ is Convergent by ACT.

NOTE: this technique helps you avoid AST here on the O.S.

2.
$$\sum_{n=1}^{\infty} (-1)^n \frac{\sin^2 n}{n^3 + 5} \to \text{Consider the A.S.} \sum_{n=1}^{\infty} \frac{\sin^2 n}{n^3 + 5}.$$

Bound the terms: $\frac{\sin^2 n}{n^3 + 5} \le \frac{\sin^2 n}{n^3} \le \frac{1}{n^3}$ and $\sum_{n=1}^{\infty} \frac{1}{n^3}$ is a convergent *p*-Series with p = 3 > 1.

Therefore, the A.S. is Convergent by CT. Finally, the O.S. is Convergent by ACT

3.
$$\sum_{n=1}^{\infty} (-1)^n \frac{n^3 + 5n}{n^8 + 7} \to \text{A.S.} \sum_{n=1}^{\infty} \frac{n^3 + 5n}{n^8 + 7} \approx \sum_{n=1}^{\infty} \frac{n^3}{n^8} = \sum_{n=1}^{\infty} \frac{1}{n^5} \text{ Conv. } p \text{-series } p = 5 > 1.$$

Check:
$$\lim_{n \to \infty} \frac{\frac{n^3 + 5n}{n^8 + 7}}{\frac{1}{n^5}} = \lim_{n \to \infty} \frac{n^8 + 5n^6}{n^8 + 7} = \lim_{n \to \infty} \frac{1 + \frac{5}{n^2}}{1 + \frac{7}{n^8}} = 1 \text{ Finite, Non-zero.}$$

Therefore the A.S. is also convergent by LCT (which by definition means that the A.S. is A.C.). Finally, the O.S. is convergent by ACT.

4. $\sum_{n=1}^{\infty} (-1)^n \frac{\arctan n}{n^2 + 9} \to \text{A.S.} \sum_{n=1}^{\infty} \frac{\arctan n}{n^2 + 9} \approx \sum_{n=1}^{\infty} \frac{\frac{\pi}{2}}{n^2} = \frac{\pi}{2} \sum_{n=1}^{\infty} \frac{1}{n^2} \text{ Constant Multiple of a Convergent } p\text{-series } p = 2 > 1 \text{ is Convergent.}$ Bound the terms: $\frac{\arctan n}{n^2 + 9} \leq \frac{\frac{\pi}{2}}{n^2 + 9} \leq \frac{\frac{\pi}{2}}{n^2}.$ We analyzed the comparison series above. Finally, the A.S. is convergent by CT which implies that the O.S. is [convergent] by ACT.

LAST NOTE: It is very important to finish the full and complete analysis of the Absolute Series. Make a conclusion about the Convergence of the Absolute Series. THEN you can use that to imply convergence of the O.S. using the ACT.