

## Functions: Fundamental Definitions and Facts

This handout should be mostly review, but some of it may be new to you. It covers much of the material in Section 7 of Saracino.

**Definition.** Let  $S, T$  be sets. A **function**  $f : S \rightarrow T$  assigns, to each  $s \in S$ , a unique  $t \in T$ .

In that case, we write  $f(s) = t$ , or  $f : s \mapsto t$ . That is,  $\boxed{\forall s \in S, \exists! t \in T \text{ s.t. } f(s) = t}$

The set  $S$  is called the **domain** of  $f$ , and the set  $T$  is called the **codomain** or **target set** of  $f$ .

(**Warning:** do *not* call  $T$  the “range” of  $T$ , as that means something different; see below.)

Actually, the above definition is a little informal. See Video 17 for a fully rigorous definition; but in practice, we will use the above informal definition, which is more intuitive.

For both definitions, the  $\exists!$  statement — i.e., “there exists a unique” — in the box above is what really matters. To clarify things in practice, if we have a rule  $f : S \rightarrow T$  which we are not yet certain is truly a function, we sometimes use the following terminology:

- To say  $f : S \rightarrow T$  is **defined** means  $\boxed{\forall s \in S, \exists t \in T \text{ s.t. } f(s) = t}$
- To say  $f : S \rightarrow T$  is **well-defined** means  $\boxed{\forall s_1, s_2 \in S, \text{ if } s_1 = s_2, \text{ then } f(s_1) = f(s_2)}$

So a function, then, is a rule/recipe  $f : S \rightarrow T$  that is **both** defined and well-defined.

**Example:**  $f : \mathbb{Z} \rightarrow \mathbb{Q}$  by  $f(x) = \frac{1}{x}$  is **not defined**

because  $0 \in \mathbb{Z}$  is in the (supposed) domain, but  $f(0) = 1/0$  doesn't make sense.

**Example:**  $f : \mathbb{Q} \rightarrow \mathbb{Z}$  by  $f\left(\frac{m}{n}\right) = m$  is **not well-defined**

because  $\frac{3}{5} = \frac{6}{10}$  but  $f\left(\frac{3}{5}\right) = 3 \neq 6 = f\left(\frac{6}{10}\right)$ .

**Definition.** Let  $S, T$  be sets, and let  $f : S \rightarrow T$  be a function.

**1** The set  $\{f(s) \mid s \in S\} = \{t \in T \mid \exists s \in S \text{ s.t. } f(s) = t\}$ , which is a subset of  $T$ , is called the **image** of  $f$  (or the **range** of  $f$ ) and is denoted  $f(S)$  or  $\text{im}(f)$ .

**2** We say  $f$  is **onto**, or **surjective**, if  $f(S) = T$ , i.e., if  $\boxed{\forall t \in T, \exists s \in S \text{ s.t. } f(s) = t}$

**3** We say  $f$  is **one-to-one**, or **injective**, if  $\boxed{\forall s_1, s_2 \in S \text{ s.t. } f(s_1) = f(s_2), \text{ we have } s_1 = s_2}$

**4** We say  $f$  is **bijective** if it is both injective and surjective.

**5** The **identity function** on  $S$  is the function  $\text{id}_S : S \rightarrow S$  given by  $\text{id}_S(x) = x$  for all  $x \in S$ .

Note that the identity function  $\text{id}_S : S \rightarrow S$ , sometimes denoted  $i_S$  or  $1_S$ , is indeed a function, and it is bijective.

Note also both the similarities **and** the differences between:

- the definitions of “ $f$  is defined” and “ $f$  is onto”
- the definitions of “ $f$  is well-defined” and “ $f$  is one-to-one.”

In particular, note that for well-defined, the implication goes one direction, but for one-to-one, it goes the other direction.

**What does it mean to say  $f_1 = f_2$ ?** It's always important in mathematics to know *precisely* what we mean when we say two objects are equal.

For functions: if  $f_1$  and  $f_2$  are functions, we say/write  $f_1 = f_2$  if:

- the domain  $S$  of  $f_1$  is the same as the domain of  $f_2$  (in the sense of equality of sets), and
- for all  $s \in S$  [i.e., for all  $s$  in this common domain], we have  $f_1(s) = f_2(s)$ .

---

**Definition.** Let  $S, T, U$  be sets, and let  $f : S \rightarrow T$  and  $g : T \rightarrow U$  be functions.

In this situation — where the domain of  $g$  is the same as the target set of  $f$  — we define the **composition** of  $g$  and  $f$ , denoted  $g \circ f$ , to be the function

$$g \circ f : S \rightarrow U \quad \text{given by} \quad g \circ f(s) = g(f(s))$$

We may read  $g \circ f$  aloud as “ $g$  composed with  $f$ ” or “ $g$  of  $f$ ” or “ $g$  after  $f$ .”

---

**Definition.** Let  $S, T$  be sets, and let  $f : S \rightarrow T$  be a function. We say  $f$  is **invertible** if there exists a function  $g : T \rightarrow S$  such that

$$f \circ g = \text{id}_T \text{ and } g \circ f = \text{id}_S$$

that is, if both:

- for all  $s \in S$ , we have  $g(f(s)) = s$ , and
- for all  $t \in T$ , we have  $f(g(t)) = t$ .

In that case,  $g$  is called the **inverse function** of  $f$ , or simply the **inverse** of  $f$ , and we write  $f^{-1} = g$ .

---

The following result is standard, but I recommend you try to prove it yourself; it is good practice both with writing if-and-only-if proofs, and with working with the various definitions in this handout. (The informal discussion on page 62 of Saracino may also be helpful.)

**Proposition.** Let  $S, T$  be sets, and let  $f : S \rightarrow T$  be a function. Then

$f$  is invertible **if and only if**  $f$  is bijective.

Moreover, in that case, the inverse function  $f^{-1} : T \rightarrow S$  is given by:

$$f^{-1}(t) = \text{the unique } s \in S \text{ such that } f(s) = t$$

for all  $t \in T$ .

---

Here's another standard result about inverses. Again, I encourage you to try to prove it yourself. (It may remind you of Theorems 3.2, 3.3, and 3.4 about groups!)

**Proposition.** Let  $S, T, U$  be sets, and let  $f : S \rightarrow T$  and  $g : T \rightarrow U$  be **invertible** functions.

**1** The inverse of  $f$  is unique. That is, if  $F : T \rightarrow S$  is a function such that  $F \circ f = \text{id}_S$  and  $f \circ F = \text{id}_T$ , then we have  $F = f^{-1}$ .

**2** The inverse function  $f^{-1}$  is also invertible, and its inverse is  $(f^{-1})^{-1} = f$ .

**3** The composition  $g \circ f$  is also invertible, and its inverse is  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .

---

Finally, composition of functions is associative; again, try to prove the following result yourself:

**Proposition.** Let  $S, T, U, V$  be sets, and let  $f : S \rightarrow T$ ,  $g : T \rightarrow U$ , and  $h : U \rightarrow V$  be functions. Then  $h \circ (g \circ f) = (h \circ g) \circ f$ .

[Note:  $h \circ (g \circ f)$  and  $(h \circ g) \circ f$  are functions from  $S$  to  $V$ . Proving they are equal means proving that for all  $s \in S$ , their values at  $s$  agree.]