

A LARGE ARBOREAL GALOIS REPRESENTATION FOR A CUBIC POSTCRITICALLY FINITE POLYNOMIAL

ROBERT L. BENEDETTO, XANDER FABER, BENJAMIN HUTZ, JAMIE JUUL,
AND YU YASUFUKU

ABSTRACT. We give a complete description of the arboreal Galois representation of a certain postcritically finite cubic polynomial over a large class of number fields and for a large class of basepoints. The associated Galois action on an infinite ternary rooted tree has Hausdorff dimension bounded strictly between that of the infinite wreath product of cyclic groups and that of the infinite wreath product of symmetric groups. We deduce a zero-density result for prime divisors in an orbit under this polynomial. We also obtain a zero-density result for the set of places of convergence of Newton's method for a certain cubic polynomial, thus resolving the first nontrivial case of a conjecture of Faber and Voloch.

1. INTRODUCTION

Let K be a field and $f \in K[z]$ a polynomial of degree $d \geq 2$. Consider Galois groups of polynomials of the form

$$f^n(z) - x, \tag{1.1}$$

where $x \in K$, and $f^n = f \circ \dots \circ f$ is the n -th iterate of f (with the convention that $f^0(z) = z$). Such groups are called *arboreal Galois groups* because (under certain hypotheses) they can be made to act on trees.

Let T_n be the graph whose vertex set is

$$\bigsqcup_{0 \leq i \leq n} f^{-i}(x),$$

and where we draw an edge from α to β if $f(\alpha) = \beta$. Let $G_n = \text{Gal}(f^n(z) - x/K)$. Clearly, G_n acts faithfully on T_n , so that $G_n \hookrightarrow \text{Aut}(T_n)$. Provided there is no critical point of f among the points of the above vertex set, the graph T_n is a regular d -ary rooted tree with root x . For such f , $\text{Aut}(T_n)$ is isomorphic to the n -fold iterated wreath product $[\mathfrak{S}_d]^n$ of the symmetric group \mathfrak{S}_d on d letters. Odoni and Juul [8, 10] showed that if $\text{char}(K)$ and the degree are not both 2, and if f is chosen generically (in the Zariski sense), then $G_n \cong \text{Aut}(T_n) \cong [\mathfrak{S}_d]^n$.

By contrast, for *specific* choices of f and x , the corresponding Galois groups may be much smaller. (See [6, §3] for a high-level explanation and [2, 4, 7, 12] for detailed examples.) Consider a polynomial f that is *postcritically finite*, or *PCF* for short, meaning that all of its critical points have finite orbit under the iteration of f . The simplest examples of PCF polynomials are the power maps $f(z) = z^d$ and the Chebyshev polynomials defined by $f(z + 1/z)^d = z^d + 1/z^d$. These two examples arise from an endomorphism of the algebraic group \mathbb{G}_m , which gives a foothold on the associated arboreal Galois representation. (A third type of example, Lattès maps, arise from an endomorphism of an elliptic curve; however, Lattès maps are never conjugate to polynomials.)

Jones and Pink [6, Thm. 3.1] have shown that for PCF maps, the Galois groups G_n have *unbounded index* inside $\text{Aut}(T_n)$ as $n \rightarrow \infty$. However, their proof does not explicitly describe G_n . One can give an upper bound for G_n inside $\text{Aut}(T_n)$ by realizing it as a specialization of $\text{Gal}((f^n(z) - t/K(t)))$, with t transcendental over K . The latter group may be embedded in the profinite monodromy group $\pi_1^{\text{ét}}(\mathbb{P}_k^1 \setminus P)$, where P is the strict postcritical orbit; this is precisely the tack taken by Pink [12] in the case of quadratic PCF polynomials.

In this paper, we give the first complete calculation of the arboreal Galois group attached to a PCF polynomial over a number field that is *not* associated with an endomorphism of an algebraic group. More specifically, we describe the Galois groups $G_n = \text{Gal}(f^n(z) - x/K)$ for the polynomial

$$f(z) = -2z^3 + 3z^2$$

over a number field K , where x is chosen to satisfy a certain local hypothesis at the primes 2 and 3. In Section 2 we will define groups E_n that fit between $\text{Aut}(T_n) \cong [\mathfrak{S}_3]^n$ and its Sylow 3-subgroup $[C_3]^n$ — the iterated wreath produce of cyclic groups of order 3. The groups E_n are somewhat tricky to handle because their action on the tree lacks a certain rigidity property: for $m < n$, the kernel of the restriction homomorphism $E_n \rightarrow E_m$ is not the direct product of copies of E_{n-m} . That is, in contrast to $[\mathfrak{S}_3]^n$ and $[C_3]^n$, the action of E_n on one branch of the tree above T_m is not independent of its action on another branch. Our main result is the following.

Theorem 1.1. *Let K be a number field, let $f(z) = -2z^3 + 3z^2 \in K[z]$, and let $x \in K$. Suppose there exist primes \mathfrak{p} and \mathfrak{q} lying over 2 and 3, respectively, such that either $v_{\mathfrak{p}}(x) = \pm 1$ or $v_{\mathfrak{p}}(1 - x) = 1$, and that $v_{\mathfrak{q}}(x) = 1$. Then for each $n \geq 1$,*

- (a) *The polynomial $f^n(z) - x$ is irreducible over K .*
- (b) *We have an isomorphism $\text{Gal}(f^n(z) - x/K) \cong E_n \subset \text{Aut}(T_n)$.*

Let $E_{\infty} = \varprojlim E_n$ and $\text{Aut}(T_{\infty}) = \varprojlim \text{Aut}(T_n)$ be the corresponding projective limits. Then the Hausdorff dimension of E_{∞} in $\text{Aut}(T_{\infty})$ is

$$\lim_{n \rightarrow \infty} \frac{\log |E_n|}{\log |\text{Aut}(T_n)|} = 1 - \frac{1}{3} \frac{\log 2}{\log 6} \approx 0.871. \quad (1.2)$$

The Galois group $\text{Gal}(f^n(z) - x/K)$ depends *a priori* on the number field K and the basepoint x , but Theorem 1.1 shows that many choices of K and x give the same isomorphism type. One key reason is that the discriminant of the second iterate is a square:

$$\text{For any } x, \quad \text{Disc}(f^2(z) - x) = [2^{16} \cdot 3^9 \cdot x^2(x-1)^2]^2. \quad (1.3)$$

This observation will be vital for forcing the Galois group of $f^n(z) - x$ to lie inside E_n .

We are also able to deduce that the geometric Galois group has the same structure:

Corollary 1.2. *Let $f(z) = -2z^3 + 3z^2$. Let t be transcendental over \mathbb{Q} . For every $n \geq 1$, we have*

$$\text{Gal}(f^n(z) - t/\bar{\mathbb{Q}}(t)) \cong E_n.$$

Odoni [11] has shown that descriptions of iterated Galois groups of this sort give rise to applications on the density of prime divisors in certain dynamically defined sequences. (See also [5, 8, 9].) More precisely, after counting elements of E_n that fix a leaf of the tree T_n , we have the following arithmetic application.

Theorem 1.3. *Let K be a number field for which there exists an unramified prime above 2 and above 3. Let $y_0 \in K \setminus \{0, 1, 3/2, -1/2\}$, and define the sequence $y_i = f^i(y_0)$. Then the set of prime ideals \mathfrak{P} such that*

$$y_i \equiv 0 \text{ or } 1 \pmod{\mathfrak{P}} \text{ for some } i \geq 0$$

has natural density zero. In particular, the set of prime divisors of the sequence (y_i) has natural density zero.

Our choice of the polynomial $f(z) = -2z^3 + 3z^2$ was originally motivated by the following conjecture of Faber and Voloch [3].

Conjecture 1.4 (Newton Approximation Fails for 100% of Primes). *Let g be a polynomial of degree $d \geq 2$ with coefficients in a number field K and let $y_0 \in K$. Define the Newton map $N(z) = z - g(z)/g'(z)$ and, for each $n \geq 0$, set $y_{n+1} = N(y_n)$. Assume the Newton approximation sequence (y_n) is not eventually periodic. Let $C(K, g, y_0)$ be the set of primes \mathfrak{P} of K for which (y_n) converges in the completion $K_{\mathfrak{P}}$ to a root of f . Then the natural density of the set $C(K, g, y_0)$ is zero.*

Faber and Voloch showed that Conjecture 1.4 holds for any polynomial g with at most 2 distinct roots. Thus, the first nontrivial case of the conjecture is a cubic polynomial with distinct roots. For reasons explained in [3, Cor. 1.2], the simplest such cubic polynomial is $g(z) = z^3 - z$, whose associated Newton map turns out to be conjugate to our polynomial $f(z) = -2z^3 + 3z^2$. Our results therefore yield a proof of the first nontrivial case of the Faber-Voloch conjecture:

Theorem 1.5. *Let K be a number field for which there exists an unramified prime over 2 and over 3. Let $g(z) = z^3 - z$. Choose $y_0 \in K$ such that the Newton iteration sequence $y_i = N^i(y_0)$ does not encounter a root of g . Then the set of primes \mathfrak{P} of K for which the Newton sequence (y_i) converges in $K_{\mathfrak{P}}$ to a root of g has natural density zero.*

The first and fourth authors, in collaboration with several others, obtained a weak form of Theorem 1.5 for a wide class of polynomials [1, Thm. 4.6]. More precisely, they showed that the density of primes as in the theorem has natural density strictly less than 1.

The outline of the paper is as follows. In Section 2, we will define and discuss the group E_n and compute the Hausdorff dimension of equation (1.2). We will then prove the rest of Theorem 1.1 in Section 3. Next, we consider the case that K is the field of rational functions $\mathbb{Q}(t)$, proving Corollary 1.2 in Section 4. In Section 5, we compute the proportion of elements of E_n that fix at least one leaf of the tree T_n , and in Section 6, we prove Theorems 1.3 and 1.5.

2. TREE AUTOMORPHISMS

Let T_n denote the regular ternary rooted tree with n levels, as in Figure 1. Note that T_n has 3^n leaves and $1 + 3 + \dots + 3^n$ vertices. Our results and many arguments will depend on an implicit labeling of the vertices of T_n . We will make this labeling explicit now for purposes of rigor, but we will not comment on it again afterward.

- The *level* of a vertex is its distance from the root.
- A vertex at level i is given a label (ℓ_1, \dots, ℓ_i) , where $\ell_j \in \{1, 2, 3\}$. The root is given the empty label $()$.
- No two vertices at the same level have the same label.




FIGURE 1. The ternary rooted tree T_n with n levels

- The unique path from the root to the vertex with label (ℓ_1, \dots, ℓ_i) consists of the vertices with labels $(\ell_1), (\ell_1, \ell_2), \dots, (\ell_1, \ell_2, \dots, \ell_i)$.

This labeling enables us to identify certain canonical subtrees of T_n . For example, for each $i \in \{1, 2, 3\}$, we consider the subtree T that is induced by the set of vertices with labels of the form $(i, *, \dots, *)$; then T is isomorphic to T_{n-1} .

The automorphism group $\text{Aut}(T_n)$ of the regular rooted ternary tree is isomorphic to the n -fold iterated wreath product $[\mathfrak{S}_3]^n$. Indeed, we may decompose T_n as a copy of T_1 (vertices of level at most 1) with 3 copies of T_{n-1} attached along the leaves of T_1 , so that

$$\text{Aut}(T_n) \cong \text{Aut}(T_{n-1}) \wr \text{Aut}(T_1) \cong [\mathfrak{S}_3]^{n-1} \wr \mathfrak{S}_3 = [\mathfrak{S}_3]^n. \quad (2.1)$$

Our labeling has the effect of fixing an isomorphism $\text{Aut}(T_1) \cong \mathfrak{S}_3$. For any elements $a_1, a_2, a_3 \in \text{Aut}(T_{n-1})$ and any $b \in \text{Aut}(T_1)$, the element

$$((a_1, a_2, a_3), b) \in \text{Aut}(T_{n-1}) \wr \text{Aut}(T_1) \cong \text{Aut}(T_n)$$

acts on the tree by first acting on the 3 copies of T_{n-1} via a_1, a_2 , and a_3 , respectively, and then by permuting these T_{n-1} 's via b . More precisely, if we label a vertex y of T_n by (x, i) , where x is a vertex of T_{n-1} and $i \in \{1, 2, 3\}$ is the vertex at the first level that y lies above, then

$$((a_1, a_2, a_3), b) \cdot (x, i) = (a_i \cdot x, b \cdot i).$$

A labeling of the leaves of T_n induces an injection $\text{Aut}(T_n) \hookrightarrow \mathfrak{S}_{3^n}$. Changing the labeling corresponds to conjugating by an element of \mathfrak{S}_{3^n} , and so each automorphism of T_n has a well defined sign attached to it, corresponding to the parity of the number of transpositions needed to represent it. Thus, for each $n \geq 1$, we have a homomorphism

$$\text{sgn}: \text{Aut}(T_n) \rightarrow \{\pm 1\}.$$

Lemma 2.1. *Let $g = ((a_1, a_2, a_3), b)$ be an element of $\text{Aut}(T_n)$ for some $n \geq 2$, where $b \in \text{Aut}(T_1)$, and $a_i \in \text{Aut}(T_{n-1})$ for $i = 1, 2, 3$. Then*

$$\text{sgn}(g) = \text{sgn}(b) \prod_{i=1}^3 \text{sgn}(a_i).$$

Proof. Partition the leaves of T_n into three disjoint sets L_1, L_2, L_3 so that the elements of L_i lie over leaf i of T_1 , for $i = 1, 2, 3$. Note that $|L_i| = 3^{n-1}$. With this notation, $\text{sgn}(a_i)$ is the sign of a_i acting as a permutation on the set L_i .

Consider first the case that $b = 1$. Then g permutes the elements of each L_i separately; hence, $\text{sgn}(g) = \prod \text{sgn}(a_i)$.

Next, consider the case that $g = ((1, 1, 1), b)$ for arbitrary $b \in \text{Aut}(T_1)$. We have already proven the desired result if $b = 1$. If b is a 2-cycle — say $b = (ij)$ — then the induced permutation on the leaves of T_n decomposes as a product of 3^{n-1} disjoint 2-cycles $(a_i a_j)$, where $a_i \in L_i$ and $a_j \in L_j$. Therefore,

$$\text{sgn}(g) = (-1)^{3^{n-1}} = -1 = \text{sgn}(b).$$

Similarly, if b is a 3-cycle, then the induced permutation on the leaves of T_n decomposes as a product of 3^{n-1} disjoint 3-cycles. Hence,

$$\text{sgn}(g) = 1 = \text{sgn}(b).$$

Finally, we consider the general case $g = ((a_1, a_2, a_3), b)$. Define $h = ((1, 1, 1), b^{-1})$. Then $hg = ((a_1, a_2, a_3), 1)$. The previous two paragraphs show that

$$\prod_{i=1}^3 \text{sgn}(a_i) = \text{sgn}(hg) = \text{sgn}(h) \text{sgn}(g) = \text{sgn}(b^{-1}) \text{sgn}(g). \quad \square$$

For any $m \leq n$, we have a restriction homomorphism $\pi_m: \text{Aut}(T_n) \rightarrow \text{Aut}(T_m)$, where T_m is the subtree with m levels with the same root vertex as T_n . We write $\text{sgn}_m = \text{sgn} \circ \pi_m$ for the composition of restriction followed by the sign map. Define a sequence of subgroups $E_n \subset \text{Aut}(T_n)$ by the following formula:

$$E_n = \begin{cases} \text{Aut}(T_1) & \text{if } n = 1 \\ (E_{n-1} \wr \text{Aut}(T_1)) \cap \ker(\text{sgn}_2) & \text{if } n \geq 2. \end{cases} \quad (2.2)$$

Here we use the embedding

$$E_{n-1} \wr \text{Aut}(T_1) \hookrightarrow \text{Aut}(T_{n-1}) \wr \text{Aut}(T_1) \cong \text{Aut}(T_n)$$

from equation (2.1). Thus, for $n \geq 2$, writing a given automorphism $\sigma \in \text{Aut}(T_n)$ as $\sigma = ((a_1, a_2, a_3), b)$, we have

$$\sigma = ((a_1, a_2, a_3), b) \in E_n \quad \text{if and only if} \quad a_1, a_2, a_3 \in E_{n-1} \text{ and } \text{sgn}_2(\sigma) = 1. \quad (2.3)$$

Proposition 2.2. *For $n \geq 1$, we have $|E_n| = 2^{3^{n-1}} \cdot 3^{\frac{3^n-1}{2}}$.*

Proof. Since $\text{Aut}(T_1) \cong \mathfrak{S}_3$, the result is clear for $n = 1$. Suppose it holds for some $n \geq 1$. Let ϕ be the composition

$$E_n \wr \text{Aut}(T_1) \hookrightarrow \text{Aut}(T_{n+1}) \xrightarrow{\text{sgn}_2} \{\pm 1\}.$$

By definition, $E_{n+1} = \ker(\phi)$, and ϕ is onto because $((1, 1, 1), \epsilon) \mapsto -1$ for any transposition ϵ of the leaves of T_1 . Thus,

$$|E_{n+1}| = \frac{1}{2} |E_n \wr \text{Aut}(T_1)| = \frac{1}{2} |\text{Aut}(T_1)| \cdot |E_n|^3 = \frac{1}{2} \cdot 6 \cdot \left(2^{3^{n-1}} \cdot 3^{\frac{3^n-1}{2}}\right)^3 = 2^{3^n} \cdot 3^{\frac{3^{n+1}-1}{2}}. \quad \square$$

Our construction of E_n depends on an identification of T_{n-1} with the subtree of T_n lying above a vertex at level 1, which in turn depends on the labeling we have assigned to T_n . In other words, E_n is not normal in $\text{Aut}(T_n)$ (for $n \geq 3$): a different labeling yields a conjugate subgroup in $\text{Aut}(T_n)$.

Proposition 2.3. E_n is normal in $\text{Aut}(T_n)$ if and only if $n = 1$ or 2 .

Proof. We have $E_1 = \text{Aut}(T_1)$, and E_2 has index 2 in $\text{Aut}(T_2)$. It remains to show that E_n is not normal in $\text{Aut}(T_n)$ for $n \geq 3$. To that end, we first construct some special elements of $\text{Aut}(T_n)$.

Define $\nu_n \in \text{Aut}(T_n)$ inductively for $n \geq 1$ as follows:

$$\nu_n = \begin{cases} (12) & n = 1 \\ ((\nu_{n-1}, 1, 1), 1) & n \geq 2. \end{cases}$$

Thus, ν_n transposes two leaves at the n -th level and acts trivially on the rest of T_n . In particular, $\text{sgn}(\nu_n) = -1$. This yields $\nu_2 \notin E_2$, and by induction, it follows that $\nu_n \notin E_n$ for $n \geq 2$. Note further that $\nu_n^{-1} = \nu_n$.

Next, for fixed $n \geq 3$, define $a = ((1, 1, 1), (123)) \in \text{Aut}(T_n)$. Then $a \in E_n$ by (2.3). However,

$$\nu_n a \nu_n^{-1} = ((\nu_{n-1}, 1, 1), 1) ((1, 1, 1), (123)) ((\nu_{n-1}, 1, 1), 1) = ((\nu_{n-1}, 1, \nu_{n-1}), (123)),$$

which does *not* belong to E_n since $\nu_{n-1} \notin E_{n-1}$ for $n \geq 3$. It follows that E_n is not normal in $\text{Aut}(T_n)$, as desired. \square

Write $T_\infty = \bigcup_{n \geq 1} T_n$ for the infinite ternary rooted tree, which has automorphism group

$$\text{Aut}(T_\infty) = \varprojlim \text{Aut}(T_n),$$

where the inverse limit is taken with respect to the restriction homomorphisms

$$\pi_m : \text{Aut}(T_n) \rightarrow \text{Aut}(T_m) \quad \text{for } m \leq n.$$

The recursive definition of E_n implies that we also have restriction homomorphisms $E_n \rightarrow E_m$ for $m \leq n$. Passing to the projective limit gives a subgroup

$$E_\infty = \varprojlim E_n$$

of $\text{Aut}(T_\infty)$.

Corollary 2.4. The Hausdorff dimension of E_∞ in $\text{Aut}(T_\infty)$ is given by equation (1.2).

Proof. Using the facts that $\text{Aut}(T_n) \cong \text{Aut}(T_{n-1}) \wr \text{Aut}(T_1)$ and that $\text{Aut}(T_1) \cong \mathfrak{S}_3$, a simple induction shows that

$$|\text{Aut}(T_n)| = 6^{\frac{3^n - 1}{2}}. \quad (2.4)$$

Combining this fact with Proposition 2.2 gives the desired result. \square

Corollary 2.5. E_∞ has infinite index in $\text{Aut}(T_\infty)$.

Comparing the cardinalities of E_n and $\text{Aut}(T_n)$, we see that they share a Sylow 3-subgroup. We can describe one such subgroup explicitly as follows. Let C_3 be the cyclic 3-subgroup of $\text{Aut}(T_1) \cong \mathfrak{S}_3$. Define a sequence of groups H_n by the following formula:

$$H_n = \begin{cases} C_3 & \text{if } n = 1 \\ H_{n-1} \wr C_3 & \text{if } n \geq 2. \end{cases}$$

We identify H_n with a subgroup of $\text{Aut}(T_n)$ using the embedding

$$H_n = H_{n-1} \wr C_3 \hookrightarrow \text{Aut}(T_{n-1}) \wr \text{Aut}(T_1) \cong \text{Aut}(T_n).$$

Evidently, $H_n \cong [C_3]^n$, the iterated wreath product. By induction, we see that

$$|H_n| = 3^{\frac{3^n - 1}{2}}, \quad (2.5)$$

so that H_n is a Sylow 3-subgroup of $\text{Aut}(T_n)$.

Proposition 2.6. *For $n \geq 1$, H_n is a Sylow 3-subgroup of E_n . It is normal in E_n if and only if $n = 1$.*

Proof. For $n = 1$, H_n is an index-2 subgroup of $E_1 = \text{Aut}(T_1)$ and, hence, it is normal. Next, for some $n \geq 1$, suppose we know that H_n is a subgroup of E_n . By their recursive definitions, to see that $H_{n+1} \subset E_{n+1}$ it suffices to show that any $h \in H_{n+1}$ restricts to an even permutation on T_2 . The restriction of h to T_2 has the form $((a_1, a_2, a_3), b)$, where each of a_1, a_2, a_3 , and b is either trivial or a 3-cycle. By Lemma 2.1, we conclude that $\text{sgn}_2(h) = 1$. Hence, $h \in E_{n+1}$, as desired.

Since $|H_n|$ is the 3-power part of $|E_n|$, we have proven the first statement of the proposition. It remains to show that H_n is not normal in E_n for $n \geq 2$.

For each $n \geq 1$, define $\tau_n \in \text{Aut}(T_n)$ inductively as follows:

$$\tau_n = \begin{cases} (12) & n = 1 \\ ((\tau_{n-1}, 1, 1), (12)) & n \geq 2. \end{cases} \quad (2.6)$$

We claim that $\tau_n \in E_n$ for all $n \geq 1$. This is clear for $n = 1$. Suppose that it holds for some $n \geq 1$. Then $((\tau_n, 1, 1), (12)) \in E_{n+1}$ if and only if its restriction to T_2 acts by an even permutation. The restriction to E_2 is given by $((12), 1, 1), (12))$, which has positive sign by Lemma 2.1.

Note that for $n \geq 2$, we have $\tau_n^{-1} = ((1, \tau_{n-1}^{-1}, 1), (12))$. Note also that for $n \geq 1$, we have $\tau_n \notin H_n$, since the restriction of τ_n to T_1 is $(12) \notin C_3$.

Next, for fixed $n \geq 2$, define $a = ((1, 1, 1), (123)) \in H_n$. Then

$$\tau_n a \tau_n^{-1} = ((\tau_{n-1}, 1, 1), (12))((1, 1, 1), (123))((1, \tau_{n-1}^{-1}, 1), (12)) = ((1, \tau_{n-1}^{-1}, \tau_{n-1}), (132)),$$

which does not belong to H_n , since $\tau_{n-1} \notin H_{n-1}$. \square

Proposition 2.7. *The Hausdorff dimension of H_∞ in $\text{Aut}(T_\infty)$ is*

$$\lim_{n \rightarrow \infty} \frac{\log |H_n|}{\log |\text{Aut}(T_n)|} = \frac{\log 3}{\log 6} \approx 0.613.$$

Proof. Immediate from equations (2.4) and (2.5). \square

Since $H_n \subseteq E_n \subseteq \text{Aut}(T_n)$, the preceding proposition and Corollary 2.4 show that for large n , E_n is substantially larger than H_n , but much smaller than $\text{Aut}(T_n)$.

Finally, we will need a lemma that constructs certain special elements of E_n :

Lemma 2.8. *Let $n \geq 2$ and let $g \in \text{Aut}(T_n)$ be an element that acts as follows:*

- *On the copy of T_{n-1} with the same root as T_n , g acts by the identity.*
- *On each copy of T_2 rooted at a vertex of T_n of level $n-2$, g acts by an even permutation of the 9 vertices.*

Then $g \in E_n$.

Proof. We proceed by induction on n . For $n = 2$, the second condition on g implies that $\text{sgn}(g) = 1$, so $g \in E_2$. Suppose that the lemma holds for $n - 1$, and let $g \in \text{Aut}(T_n)$ satisfy the given conditions. Let u_1, u_2, u_3 be the vertices of T_n at level 1. Write $T_{n-1}(u_i)$ for the copy of T_{n-1} inside T_n that is rooted at u_i . Then g restricts to an element of $\text{Aut}(T_{n-1}(u_i))$ that satisfies the two conditions of the lemma. By the induction hypothesis, $g|_{T_{n-1}(u_i)} \in E_{n-1}$ for $i = 1, 2, 3$. In addition, g is the identity and, hence, is even, on T_2 . Thus, $g \in E_n$ by the criterion of (2.3). \square

3. MAIN THEOREM

Let K be a field of characteristic 0, and consider the polynomial

$$f(z) = -2z^3 + 3z^2 \in K[z].$$

The critical points of f in \mathbb{P}^1 are 0, 1, and ∞ , all of which are fixed by f . Hence, f is PCF and, in fact, the union of the forward orbits of its critical points is $\{0, 1, \infty\}$. Choose a point $x \in K \setminus \{0, 1\}$ to be the root of our preimage tree. Note that there is no critical point in the backward orbit of x .

For each $n \geq 1$, define

$$K_n = K(f^{-n}(x)) \quad \text{and} \quad G_n = \text{Gal}(K_n/K). \quad (3.1)$$

Lemma 3.1. *For any field K of characteristic zero and any $x \in K \setminus \{0, 1\}$, the Galois group G_n of (3.1) is isomorphic to a subgroup of E_n .*

Proof. Because there is no critical point in the backward orbit of x , the set $f^{-i}(x)$ consists of exactly 3^i distinct elements for each $i \geq 0$. Identify the vertices of the ternary tree T_n with the set $\bigsqcup_{0 \leq i \leq n} f^{-i}(x)$, with vertex y lying immediately above y' if and only if $f(y) = y'$. This identification induces a faithful action of G_n on T_n and, hence, G_n may be identified with a subgroup of $\text{Aut}(T_n)$.

To see that this subgroup G_n lies inside E_n , we proceed by induction on n . For $n = 1$, this is clear because $E_1 = \text{Aut}(T_1)$.

Fix $n \geq 2$, and assume we know the lemma holds for $n - 1$. Write $f^{-1}(x) = \{y_1, y_2, y_3\}$. For each $i = 1, 2, 3$, applying the lemma to the field $K(y_i)$ with root point y_i shows that

$$\text{Gal}\left(K(f^{-(n-1)}(y_i))/K(y_i)\right)$$

is a subgroup of E_{n-1} . (The labeling of T_n allows us to identify the portion of T_n above y_i with T_{n-1} .) Since $\text{Gal}(K_1/K)$ is a subgroup of $\text{Aut}(T_1)$, it follows that G_n is isomorphic to a subgroup B_n of $E_{n-1} \wr \text{Aut}(T_1)$.

It remains to show that $B_n \subseteq \ker(\text{sgn}_2)$. Direct computation shows that the discriminant of the degree-nine polynomial $f^2(z) - x$ is given by equation (1.3). Since this discriminant is a square in K , all elements of B_n act as even permutations of the nine points of $f^{-2}(x)$. Thus, $B_n \subseteq \ker(\text{sgn}_2)$, as desired. \square

Our goal is to compute the arboreal Galois groups G_n in the case that K is a number field and that the basepoint $x \in K \setminus \{0, 1\}$ satisfies the following local hypothesis:

There exist primes \mathfrak{p} and \mathfrak{q} of K lying above 2 and 3, respectively, (†)
such that $v_{\mathfrak{q}}(x) = 1$, and either $v_{\mathfrak{p}}(x) = \pm 1$ or $v_{\mathfrak{p}}(1 - x) = 1$.

If this hypothesis holds, we will say that the pair (K, x) satisfies property (\dagger) (relative to \mathfrak{p} and \mathfrak{q}).

Example 3.2. If $K = \mathbb{Q}$, then the pairs $(\mathbb{Q}, 3)$ and $(\mathbb{Q}, 3/2)$ both satisfy property (\dagger) . The latter pair will be important for our arithmetic applications.

Lemma 3.3. *Suppose that (K, x) satisfies (\dagger) relative to \mathfrak{p} and \mathfrak{q} . Then $f^n(z) - x$ is Eisenstein at \mathfrak{q} for all $n \geq 1$. In particular, $f^n(z) - x$ is irreducible for all $n \geq 1$.*

Proof. A simple induction shows that $f^n(z) \in \mathbb{Z}[z]$ has degree 3^n with constant term 0 and leading coefficient $c = (-2)^{(3^{n+1}-1)/2}$. Thus, all coefficients a of $f^n(z) - x$ satisfy $v_{\mathfrak{q}}(a) \geq 0$, with lead coefficient c satisfying $v_{\mathfrak{q}}(c) = 0$, and with constant term x satisfying $v_{\mathfrak{q}}(x) = 1$. \square

Proposition 3.4. *Let K be a number field, and let $x \in K$. Suppose that (K, x) satisfies property (\dagger) relative to primes \mathfrak{p} and \mathfrak{q} . Let $n \geq 0$, and let $y \in f^{-n}(x)$. Then:*

- (1) *There are primes \mathfrak{p}' and \mathfrak{q}' of $K(y)$ lying above \mathfrak{p} and \mathfrak{q} , respectively, such that*

$$e(\mathfrak{p}'/\mathfrak{p}) = 2^n \quad \text{and} \quad e(\mathfrak{q}'/\mathfrak{q}) = 3^n.$$

- (2) *The pair $(K(y), y)$ satisfies property (\dagger) relative to \mathfrak{p}' and \mathfrak{q}' .*

Proof. We proceed by induction on n . The statement is trivial for $n = 0$. We therefore assume for the rest of the proof that $n \geq 1$ and that the statement holds for $n - 1$.

Given $y \in f^{-n}(x)$, let $y'' = f(y) \in f^{-(n-1)}(x)$. By our inductive hypothesis, there are primes \mathfrak{p}'' and \mathfrak{q}'' of $K(y'')$ lying over \mathfrak{p} and \mathfrak{q} satisfying the desired properties for $n - 1$. The polynomial

$$f(z) - y'' = -2z^3 + 3z^2 - y'' \in K(y'')[z]$$

is Eisenstein at \mathfrak{q}'' . Thus, there is only one prime \mathfrak{q}' of $K(y)$ lying above \mathfrak{q}'' , with ramification index $e(\mathfrak{q}'/\mathfrak{q}'') = 3$, and with $v_{\mathfrak{q}'}(y) = 1$. Moreover,

$$e(\mathfrak{q}'/\mathfrak{q}) = e(\mathfrak{q}'/\mathfrak{q}'') \cdot e(\mathfrak{q}''/\mathfrak{q}) = 3 \cdot 3^{n-1} = 3^n.$$

Meanwhile, by statement (2) for $n - 1$, we have either $v_{\mathfrak{p}''}(y'') = 1$, $v_{\mathfrak{p}''}(1 - y'') = 1$, or $v_{\mathfrak{p}''}(y'') = -1$. We consider these three cases separately.

If $v_{\mathfrak{p}''}(y'') = 1$, then the Newton polygon of $f(z) - y''$ at \mathfrak{p}'' has a segment of length 2 and height 1. Thus, there is a prime \mathfrak{p}' of $K(y)$ lying above \mathfrak{p}'' , with ramification index $e(\mathfrak{p}'/\mathfrak{p}'') = 2$, and with $v_{\mathfrak{p}'}(y) = 1$. Moreover,

$$e(\mathfrak{p}'/\mathfrak{p}) = e(\mathfrak{p}'/\mathfrak{p}'') \cdot e(\mathfrak{p}''/\mathfrak{p}) = 2 \cdot 2^{n-1} = 2^n.$$

If $v_{\mathfrak{p}''}(1 - y'') = 1$, note that f is self-conjugate via $z \mapsto 1 - z$; that is, $1 - f(1 - z) = f(z)$. Thus, $1 - y \in f^{-1}(1 - y'')$, and the previous paragraph applied to $1 - y$ gives the desired conclusion.

Finally, if $v_{\mathfrak{p}''}(y'') = -1$, then because $v_{\mathfrak{p}''}(-2) \geq 1$, the Newton polygon of $f(z) - y''$ at \mathfrak{p}'' has a segment of length 2 and height -1 . Thus, there is a prime \mathfrak{p}' of $K(y)$ lying above \mathfrak{p}'' , with ramification index $e(\mathfrak{p}'/\mathfrak{p}'') = 2$, and with $v_{\mathfrak{p}'}(y) = -1$. Once again, then, we have $e(\mathfrak{p}'/\mathfrak{p}) = 2^n$. \square

Corollary 3.5. *Let K and x be as in Proposition 3.4. Let $n \geq 1$ and let K_n be the splitting field of $f^n(z) - x$ over K . Then*

$$6^n \mid [K_n : K].$$

Proof. Pick $y \in f^{-n}(x)$, and let \mathfrak{p}' and \mathfrak{q}' be the primes of $K(y)$ given by Proposition 3.4. Since K_n/K has intermediate extension $K(y)/K$, the ramification index of some prime of K_n over \mathfrak{p} must be divisible by $e(\mathfrak{p}'/\mathfrak{p}) = 2^n$. Similarly, the ramification index of some prime of K_n over \mathfrak{q} must be divisible by $e(\mathfrak{q}'/\mathfrak{q}) = 3^n$. Thus, $6^n \mid [K_n : K]$. \square

Proposition 3.6. *Let K and x be as in Proposition 3.4. Then*

$$\text{Gal}\left(K(f^{-1}(x))/K\right) \cong E_1 \cong \mathfrak{S}_3 \quad \text{and} \quad \text{Gal}\left(K(f^{-2}(x))/K\right) \cong E_2.$$

Proof. Let $K_1 = K(f^{-1}(x))$ and $K_2 = K(f^{-2}(x))$. Then, as a splitting field of a cubic polynomial, K_1 is Galois over K with $\text{Gal}(K_1/K)$ isomorphic to a subgroup of \mathfrak{S}_3 . By Corollary 3.5 with $n = 1$, we have $6 \mid [K_1 : K]$. Hence, $\text{Gal}(K_1/K) \cong \mathfrak{S}_3$.

By Lemma 3.1, $\text{Gal}(K_2/K)$ acts on $f^{-2}(x)$ as a subgroup of E_2 . It suffices to show that every element of E_2 is realized in $\text{Gal}(K_2/K)$.

Write $f^{-1}(x) = \{u_1, u_2, u_3\}$, and for each $i = 1, 2, 3$, write

$$f^{-1}(u_i) = \{v_{i1}, v_{i2}, v_{i3}\}.$$

Claim 1. There exists $\tau \in \text{Gal}(K_2/K_1) \subseteq \text{Gal}(K_2/K)$ that

- fixes v_{1j} for each $j = 1, 2, 3$,
- acts as a 2-cycle on the set $f^{-1}(u_2) = \{v_{21}, v_{22}, v_{23}\}$, and
- acts as a 2-cycle on the set $f^{-1}(u_3) = \{v_{31}, v_{32}, v_{33}\}$.

To prove Claim 1, note that $36 \mid [K_2 : K]$ by Corollary 3.5 and, hence, $6 \mid [K_2 : K_1]$, since $[K_1 : K] = 6$. By Cauchy's Theorem, there is some $\tau_1 \in \text{Gal}(K_2/K_1)$ of order 2. Since τ_1 fixes each u_i , it must act on each set $f^{-1}(u_i)$ as an element of \mathfrak{S}_3 of order dividing 2. Thus, for each $i = 1, 2, 3$, τ_1 acts as either a 2-cycle or the identity on $f^{-1}(u_i)$. In addition, τ_1 is even as a permutation on $f^{-2}(x)$ but (being of order 2) is not the identity. Thus, τ_1 acts as a 2-cycle on the preimages of exactly two of u_1, u_2, u_3 , and as the identity on the preimage of the third, u_m .

Choose $\gamma' \in \text{Gal}(K_1/K)$ with $\gamma'(u_1) = u_m$, and lift γ' to $\gamma \in \text{Gal}(K_2/K)$. Let $\tau = \gamma^{-1}\tau_1\gamma$. Then $\tau(u_i) = u_i$ for each i , and τ satisfies each of the three bulleted properties, proving Claim 1.

Claim 2. There exists $\rho \in \text{Gal}(K_2/K_1) \subseteq \text{Gal}(K_2/K)$ that acts as

- a 3-cycle on $f^{-1}(u_1) = \{v_{11}, v_{12}, v_{13}\}$,
- either a 3-cycle or the identity on $f^{-1}(u_2) = \{v_{21}, v_{22}, v_{23}\}$, and
- either a 3-cycle or the identity on $f^{-1}(u_3) = \{v_{31}, v_{32}, v_{33}\}$.

To prove Claim 2, we again note that $6 \mid [K_2 : K_1]$. Hence, by Cauchy's Theorem, there is some $\rho_1 \in \text{Gal}(K_2/K_1)$ of order 3. We have $\rho_1(u_i) = u_i$ for each i , with ρ_1 acting as a 3-cycle on either one, two, or all three of $f^{-1}(u_i)$, and as the identity on the others. Lifting some appropriate $\gamma' \in \text{Gal}(K_1/K)$ to $\gamma \in \text{Gal}(K_2/K)$, we can ensure that $\rho = \gamma^{-1}\rho_1\gamma$ satisfies the desired properties, proving Claim 2.

By the conclusions of Claims 1 and 2, the permutation $\tau\rho$ acts as

- a 3-cycle on $f^{-1}(u_1) = \{v_{11}, v_{12}, v_{13}\}$,
- a 2-cycle on $f^{-1}(u_2) = \{v_{21}, v_{22}, v_{23}\}$, and
- a 2-cycle on $f^{-1}(u_3) = \{v_{31}, v_{32}, v_{33}\}$.

Let $\sigma = (\tau\rho)^2$. Then σ acts as a 3-cycle on $f^{-1}(u_1)$ and as the identity on $f^{-1}(u_2) \cup f^{-1}(u_3)$.

Conjugating σ by permutations γ as in the proof of Claim 1 and then composing the resulting conjugates with one another, we see that $\text{Gal}(K_2/K_1)$ contains each element of E_2 that is the identity on $f^{-1}(x)$ and is either the identity or a 3-cycle on each $f^{-1}(u_i)$. There are $3^3 = 27$ such permutations.

Conjugating τ by permutations from the previous paragraph, as well as by permutations γ as in the proof of Claim 1, we also see that $\text{Gal}(K_2/K_1)$ contains each element of E_2 that is the identity on $f^{-1}(x)$, is a 2-cycle on exactly two of $f^{-1}(u_1)$, $f^{-1}(u_2)$, and $f^{-1}(u_3)$, and is the identity on the third. There are $3^3 = 27$ such permutations.

Composing the maps of the previous two paragraphs, we see that $\text{Gal}(K_2/K_1)$ contains each element of E_2 that is the identity on $f^{-1}(x)$, is a 2-cycle on exactly two of $f^{-1}(u_1)$, $f^{-1}(u_2)$, and $f^{-1}(u_3)$, and is a 3-cycle on the third. There are $3^3 \cdot 2 = 54$ such permutations.

Thus, $[K_2 : K_1] \geq 27 + 27 + 54 = 108$, and, hence,

$$[K_2 : K] \geq 6 \cdot 108 = 648 = |E_2|.$$

Since $\text{Gal}(K_2/K)$ is isomorphic to a subgroup of E_2 , we must have $\text{Gal}(K_2/K) \cong E_2$. \square

We are now prepared to prove part (b) of Theorem 1.1, which we restate now.

Theorem 3.7. *Let K and x be as in Proposition 3.4. Then for any $n \geq 1$,*

$$\text{Gal}\left(K(f^{-n}(x))/K\right) \cong E_n.$$

Proof. By Lemma 3.1, we know that the Galois group is isomorphic to a subgroup of E_n . We must show that this subgroup is E_n itself. We proceed by induction on n . For $n = 1, 2$, we are done by Proposition 3.6. Assuming we know the statement (for any such K and x) for a particular $n \geq 2$, we will now show it for $n + 1$.

Let

$$K_n = K(f^{-n}(x)) \quad \text{and} \quad K_{n+1} = K(f^{-(n+1)}(x)).$$

Write $f^{-1}(x) = \{u_1, u_2, u_3\}$. By Proposition 3.4, each of the pairs $(K(u_i), u_i)$ satisfies property (\dagger) for $i = 1, 2, 3$. Thus, our induction hypothesis says that $\text{Gal}(K_n/K)$ and all three of the Galois groups

$$\text{Gal}\left(K(f^{-n}(u_i))/K(u_i)\right), \quad \text{for } i = 1, 2, 3$$

are isomorphic to E_n . Pick

$$y \in f^{-(n-2)}(u_3) \subseteq f^{-(n-1)}(x).$$

See Figure 2. Our main goal, which we will achieve at the end of Step 3 below, is to construct an element $\lambda \in \text{Gal}(K_{n+1}/K)$ that is the identity on

$$f^{-(n+1)}(x) \setminus f^{-2}(y) \quad \text{and on} \quad f^{-1}(y),$$

but which acts as two disjoint 2-cycles on $f^{-2}(y)$.

Step 1. Define

$$H = \text{Gal}\left(K(f^{-n}(u_3))/K(u_3)\right).$$

By the induction hypothesis, $H \cong E_n$. By Lemma 2.8 there is some $\sigma_1 \in H$ that is the identity on

$$f^{-n}(u_3) \setminus f^{-2}(y) \quad \text{and on} \quad f^{-1}(y)$$




FIGURE 2. The locations of x , u_1 , u_2 , u_3 and y in T_{n+1} .

and acts as two 2-cycles on $f^{-2}(y)$. Lift σ_1 to

$$\sigma \in \text{Gal}(K_{n+1}/K(u_3)) \subseteq \text{Gal}(K_{n+1}/K).$$

Thus, σ acts as we would like λ to act on $f^{-n}(u_3)$, but we have no idea how it acts on $f^{-n}(u_1)$ and $f^{-n}(u_2)$.

Step 2. By Lemma 2.8, we may pick $\tau_1 \in \text{Gal}(K_n/K) \cong E_n$ that is the identity on

$$f^{-n}(x) \setminus f^{-1}(y)$$

and acts as a 3-cycle on $f^{-1}(y)$. Lift τ_1 to $\tau \in \text{Gal}(K_{n+1}/K)$.

Then $\tau\sigma\tau^{-1} \in \text{Gal}(K_{n+1}/K)$ acts as

- the identity on $f^{-n}(u_3) \setminus f^{-2}(y)$ and $f^{-1}(y)$,
- two 2-cycles on $f^{-2}(y)$, and
- the same as σ on $f^{-(n-1)}(\{u_1, u_2\})$,

where the two 2-cycles on $f^{-2}(y)$ for $\tau\sigma\tau^{-1}$ do not occur above the same two elements of $f^{-1}(y)$ as the two 2-cycles for σ .

Thus, $\tau\sigma\tau^{-1}\sigma^{-1} \in \text{Gal}(K_{n+1}/K)$ acts as

- the identity on $f^{-n}(u_3) \setminus f^{-2}(y)$ and $f^{-1}(y)$,
- two 2-cycles and (perhaps) a separate 3-cycle on $f^{-2}(y)$, and
- the identity on $f^{-(n-1)}(\{u_1, u_2\})$.

Cubing to kill the possible 3-cycle in $f^{-2}(y)$, we see that

$$\rho = (\tau\sigma\tau^{-1}\sigma^{-1})^3 \in \text{Gal}(K_{n+1}/K)$$

acts as

- the identity on $f^{-n}(x)$,
- the identity on $f^{-n}(u_3) \setminus f^{-2}(y)$, and
- two 2-cycles on $f^{-2}(y)$.

Step 3. Consider the permutations $\tau, \rho \in \text{Gal}(K_{n+1}/K)$ of Step 2. Then $\tau\rho\tau^{-1}\rho^{-1}$ acts as

- the identity on $f^{-n}(x)$,
- the identity on $f^{-n}(u_3) \setminus f^{-2}(y)$,
- two 2-cycles and (perhaps) a separate 3-cycle on $f^{-2}(y)$, and
- for each $v \in f^{-(n-1)}(\{u_1, u_2\})$, an even permutation of $f^{-1}(v)$.

The even permutations of the last bullet point above are even permutations in \mathfrak{S}_3 , and, hence, each is either the identity or a 3-cycle. Cubing, we see that

$$\lambda = (\tau\rho\tau^{-1}\rho^{-1})^3 \in \text{Gal}(K_{n+1}/K)$$

acts as the identity on

$$f^{-(n+1)}(x) \setminus f^{-2}(y) \quad \text{and on} \quad f^{-1}(y),$$

and it acts as two 2-cycles on $f^{-2}(y)$, achieving the main goal from the start of the proof.

Step 4. Recall $H = \text{Gal}(K(f^{-n}(u_3))/K(u_3)) \cong E_n$. Pick $w \in f^{-1}(y)$. By Lemma 2.8, we can pick $\gamma_1 \in H$ that is the identity on

$$f^{-n}(u_3) \setminus f^{-1}(w)$$

and acts as a 3-cycle on $f^{-1}(w)$. Lift γ_1 to $\gamma \in \text{Gal}(K_{n+1}/K)$.

Conjugating the permutation λ (of Step 3) by various products of γ and the permutation τ (of Step 2), we see that $\text{Gal}(K_{n+1}/K)$ contains each of the 27 permutations that is the identity on

$$Y = (f^{-(n+1)}(x) \setminus f^{-2}(y)) \cup f^{-1}(y)$$

and acts as two disjoint 2-cycles on $f^{-2}(y)$. In addition, taking products of pairs of such permutations, $\text{Gal}(K_{n+1}/K)$ contains all 27 permutations that are the identity on Y and products of disjoint 3-cycles on $f^{-2}(y)$. Still taking products of pairs, $\text{Gal}(K_{n+1}/K)$ also contains all 54 permutations that are the identity on Y and the product of a disjoint 3-cycle and two 2-cycles on $f^{-2}(y)$. Together, then, $\text{Gal}(K_{n+1}/K)$ contains a subgroup H_y that acts trivially on Y , with $|H_y| = 108 = 2^2 \cdot 3^3$.

Since $f^{n-1}(z) - x$ is irreducible over K , for each root $y' \in f^{-(n-1)}(x)$ there is some

$$\delta_1 \in \text{Gal}(K_{n-1}/K) \cong E_{n-1}$$

with $\delta_1(y') = y$. Lift δ_1 to $\delta \in \text{Gal}(K_{n+1}/K)$. Then,

$$H_{y'} = \delta^{-1}H_y\delta$$

is a 108-element subgroup of $\text{Gal}(K_{n+1}/K)$ that acts trivially on

$$Y' = (f^{-(n+1)}(x) \setminus f^{-2}(y')) \cup f^{-1}(y').$$

There are 3^{n-1} choices for y' , and any two of the resulting subgroups $H_{y'}$ act nontrivially on disjoint portions of the preimage tree. In addition, they all act trivially on $f^{-n}(x)$ and, hence, trivially on K_n . Thus, the product of all of them forms a subgroup $B \subseteq \text{Gal}(K_{n+1}/K_n)$ of order

$$(2^2 \cdot 3^3)^{3^{(n-1)}} = \frac{|E_{n+1}|}{|E_n|},$$

by Proposition 2.2. Hence,

$$|\text{Gal}(K_{n+1}/K)| = |\text{Gal}(K_{n+1}/K_n)| \cdot |\text{Gal}(K_n/K)| \geq |B| \cdot |E_n| = |E_{n+1}|.$$

Since $\text{Gal}(K_{n+1}/K)$ is isomorphic to a subgroup of the finite group E_{n+1} , we must therefore have $\text{Gal}(K_{n+1}/K) \cong E_{n+1}$. \square

4. THE GEOMETRIC REPRESENTATION

Let L be a number field, and consider the rational function field $K = L(t)$. Since $t \in L(t) \setminus \{0, 1\}$, Lemma 3.1 states that the Galois group of $L(f^{-n}(t))$ over $L(t)$ is a subgroup of E_n . In fact, we have the following much stronger statement.

Proposition 4.1. *Let L be a number field and t a transcendental element over L . Then*

$$\text{Gal}\left(L(f^{-n}(t))/L(t)\right) \cong E_n.$$

Proof. Let $G_n = \text{Gal}(L(f^{-n}(t))/L(t))$. We can choose $x \in L$ such that the pair (L, x) satisfies property (\dagger) of Section 3. By Theorem 3.7, we have

$$\text{Gal}\left(L(f^{-n}(x))/L(x)\right) = \text{Gal}\left(L(f^{-n}(x))/L\right) \cong E_n.$$

Therefore, the specialization lemma of [10, Lem. 2.4] implies that G_n has a subgroup isomorphic to E_n . On the other hand, applying Lemma 3.1 to G_n shows that G_n is isomorphic to a subgroup of E_n . Since E_n is finite, G_n must be isomorphic to E_n . \square

Corollary 4.2. $\text{Gal}\left(\bar{\mathbb{Q}}(f^{-n}(t))/\bar{\mathbb{Q}}(t)\right) \cong E_n$.

Proof. Since the previous proposition holds for any number field, \mathbb{Q} must be algebraically closed in $\mathbb{Q}(f^{-n}(t))$. Hence,

$$\text{Gal}\left(\bar{\mathbb{Q}}(f^{-n}(t))/\bar{\mathbb{Q}}(t)\right) \cong \text{Gal}\left(\mathbb{Q}(f^{-n}(t))/\mathbb{Q}(t)\right) \cong E_n. \quad \square$$

5. COUNTING ELEMENTS THAT FIX LEAVES OF T_n

Write $E_{n,\text{fix}}$ for the set of elements of E_n that fix at least one leaf of T_n . The goal of this section is to show that $E_{n,\text{fix}}$ constitutes a vanishingly small proportion of E_n :

Theorem 5.1. *We have*

$$\lim_{n \rightarrow \infty} \frac{|E_{n,\text{fix}}|}{|E_n|} = 0.$$

We begin by finding a recursive formula for $|E_{n,\text{fix}}|$ in terms of certain auxiliary quantities. For $n \geq 1$ and $i \in \{1, 2, 3\}$, we define the following:

$$\begin{aligned} A_{n,i} &= \left| \{s \in E_n \mid s \text{ acts as an element of order } i \text{ on } T_1\} \right| \\ A'_{n,i} &= \left| \{s \in E_n \mid s \text{ acts as an element of order } i \text{ on } T_1 \text{ and fixes a leaf of } T_n\} \right|. \end{aligned}$$

For example, if $n = 1$, we have

$$\begin{aligned} A_{1,1} &= 1 & A'_{1,1} &= 1 \\ A_{1,2} &= 3 & A'_{1,2} &= 3 \\ A_{1,3} &= 2 & A'_{1,3} &= 0. \end{aligned}$$

For any $n \geq 1$, note that $A'_{n,3} = 0$, because an element s that permutes the leaves of T_1 by a 3-cycle cannot fix a leaf of T_n . It follows that

$$|E_{n,\text{fix}}| = A'_{n,1} + A'_{n,2} \quad \text{and} \quad |E_n| = A_{n,1} + A_{n,2} + A_{n,3}.$$

Lemma 5.2. For $n \geq 1$, we have

$$A_{n,2} = 3A_{n,1}, \quad A_{n,3} = 2A_{n,1}, \quad \text{and} \quad |E_n| = 6A_{n,1}.$$

Proof. The restriction homomorphism $\pi: E_n \rightarrow E_1$ is surjective. Indeed, we see that $\pi((1, 1, 1), (123)) = (123)$, and $\pi(\tau_n) = (12)$, where τ was defined in (2.6).

Now $A_{n,1} = |\ker(\pi)|$. The result then follows because E_1 contains one element of order 1, three elements of order 2, and two elements of order 3. \square

Lemma 5.3. For $n \geq 1$, we have

$$|E_{n+1}| = 3|E_n|^3.$$

Proof. Apply Proposition 2.2. \square

Lemma 5.4. For $n \geq 1$, we have

$$A'_{n+1,1} = 54A_{n,1}^2 (A'_{n,1} + A'_{n,2}) - 9A_{n,1} (A'_{n,1} + A'_{n,2})^2 + 3A'_{n,1} (A'_{n,2})^2 + (A'_{n,1})^3.$$

Proof. Let $s \in E_{n+1}$ be an element that acts as the identity on T_1 . Then the restriction of s to T_2 is of the form $((a_1, a_2, a_3), 1)$ for some $a_1, a_2, a_3 \in \text{Aut}(T_1)$. By Lemma 2.1, the fact that this element lies in E_2 means $1 = \prod \text{sgn}(a_i)$. So among the a_i 's, there are either zero 2-cycles or exactly two 2-cycles. We treat these cases separately.

Case 1: Zero 2-cycles. As an element of E_{n+1} , we have $s = ((\tilde{a}_1, \tilde{a}_2, \tilde{a}_3), 1)$, where each $\tilde{a}_i \in E_n$ restricts to either the identity or a 3-cycle on T_1 . The total number of elements \tilde{a}_i of this shape is $A_{n,1} + A_{n,3}$, while the number that do not fix a leaf of T_n is $(A_{n,1} + A_{n,3} - A'_{n,1})$. Thus, the number of elements of $E_{n+1, \text{fix}}$ that act as the identity on T_1 but with no 2-cycle on T_2 is

$$(A_{n,1} + A_{n,3})^3 - (A_{n,1} + A_{n,3} - A'_{n,1})^3 = 27A_{n,1}^2 A'_{n,1} - 9A_{n,1} (A'_{n,1})^2 + (A'_{n,1})^3, \quad (5.1)$$

where we applied Lemma 5.2 when we expanded the two expressions.

Case 2: Two 2-cycles. As an element of E_{n+1} , we have $s = ((\tilde{a}_1, \tilde{a}_2, \tilde{a}_3), 1)$, where two of the $\tilde{a}_i \in E_n$ restrict to 2-cycles on T_1 , and the remaining \tilde{a}_i restricts to the identity or a 3-cycle. There are 3 choices for the index i_0 such that \tilde{a}_{i_0} is the identity or a 3-cycle. For a given choice of i_0 , there are

$$(A'_{n,1} + A'_{n,3})A_{n,2}^2 = A'_{n,1}A_{n,2}^2 = 9A_{n,1}^2 A'_{n,1}$$

choices of triples $(\tilde{a}_1, \tilde{a}_2, \tilde{a}_3)$ such that s fixes a leaf of the copy of T_n above the i_0 -leaf of T_1 .

For s not to fix a leaf of the T_n above the i_0 -leaf, at least one of the other two \tilde{a}_i 's (each of which acts as a 2-cycle on T_1) must fix a leaf of T_n . By inclusion-exclusion, the number of choices for this pair of \tilde{a}_i 's is

$$2A_{n,2}A'_{n,2} - (A'_{n,2})^2 = (6A_{n,1} - A'_{n,2}) A'_{n,2}.$$

For $i = i_0$, the element $\tilde{a}_i \in E_n$ acts as the identity or as a 3-cycle on T_1 but fixes no leaf of T_n . The number of such elements of E_n is

$$A_{n,1} + A_{n,3} - A'_{n,1} = 3A_{n,1} - A'_{n,1}.$$

(Again, we have applied Lemma 5.2 in all three displayed equations above.)

Thus, the number of elements $s \in E_{n+1, \text{fix}}$ that act as the identity on T_1 and as two 2-cycles on the leaves of T_2 is

$$3 \left[9A_{n,1}^2 A'_{n,1} + (3A_{n,1} - A'_{n,1}) (6A_{n,1} - A'_{n,2}) A'_{n,2} \right]. \quad (5.2)$$

Adding (5.1) and (5.2) and expanding yields the desired result. \square

Lemma 5.5. *For $n \geq 1$, we have*

$$A'_{n+1,2} = 54A_{n,1}^2 (A'_{n,1} + A'_{n,2}).$$

Proof. Let $s \in E_{n+1}$ be an element that acts as a 2-cycle on T_1 . Then the restriction of s to T_2 is of the form $((a_1, a_2, a_3), b)$, where $a_1, a_2, a_3, b \in \text{Aut}(T_1)$, and where b is a 2-cycle. There are 3 choices for b , and each will give the same count. Without loss, then, let us suppose that $b = (12)$. By Lemma 2.1, we have $\prod \text{sgn}(a_i) = -1$. Thus, either exactly one or all three of the a_i are 2-cycles.

Case 1: a_3 is a 2-cycle. Then $\text{sgn}(a_1)\text{sgn}(a_2) = 1$. Write $s \in E_{n+1}$ as $s = ((\tilde{a}_1, \tilde{a}_2, \tilde{a}_3), b)$, with $\tilde{a}_i \in E_n$. Since the fixed point of s cannot lie above the 1- or 2-branch of T_1 , there are

$$A_{n,2}^2 + (A_{n,1} + A_{n,3})^2 = 18A_{n,1}^2$$

choices for the pair $(\tilde{a}_1, \tilde{a}_2)$, where we have applied Lemma 5.2. Now \tilde{a}_3 must fix a leaf of T_n , and hence there are $A'_{n,2}$ choices for \tilde{a}_3 . Thus, the total number of such elements s is

$$18A_{n,1}^2 A'_{n,2}. \quad (5.3)$$

Case 2: a_3 is not a 2-cycle. Then $\text{sgn}(a_1)\text{sgn}(a_2) = -1$. Write $s = ((\tilde{a}_1, \tilde{a}_2, \tilde{a}_3), b)$ as before. Exactly one of a_1 or a_2 is a 2-cycle, and since the fixed point of s cannot lie above the 1- or 2-branch of T_1 , there are

$$2A_{n,2}(A_{n,1} + A_{n,3}) = 18A_{n,1}^2$$

choices for the pair $(\tilde{a}_1, \tilde{a}_2)$. Now \tilde{a}_3 must fix a leaf of T_n , and hence it must restrict to the identity on T_1 , yielding $A'_{n,1}$ options for \tilde{a}_3 . Thus, the total number of such elements s is

$$18A_{n,1}^2 A'_{n,1}. \quad (5.4)$$

Adding (5.3) and (5.4) and multiplying by 3 (the number of choices for the 2-cycle b) yields the desired result. \square

Lemma 5.6. *Let $\phi(t) = t - \frac{1}{2}t^2 + \frac{1}{3}t^3$. Then ϕ is increasing on $(0, 1)$, and for any $x \in [0, 1]$, we have $\phi^n(x) \rightarrow 0$ as $n \rightarrow \infty$.*

Proof. The first statement is evident from looking at the derivative. For the second, we claim that $0 \leq \phi(x) \leq x$ for all $x \in [0, 1]$. Indeed, this follows from

$$\phi(x) = x - \frac{1}{2}x^2 \left(1 - \frac{2}{3}x \right) \leq x,$$

and

$$\phi(x) = \frac{x}{3} \left(x^2 - \frac{3}{2}x + 3 \right) = \frac{x}{3} \left[\left(x - \frac{3}{4} \right)^2 + \frac{39}{16} \right] \geq 0.$$

Thus, defining $x_n = \phi^n(x)$ for $n \geq 0$, we see that the sequence $\{x_n\}_{n \geq 0}$ is a nonincreasing sequence in $[0, 1]$ and, hence, has a limit $y \in [0, 1]$. By continuity, we must have $\phi(y) = y$. However, $\phi(y) - y = 0$ yields $y = 0$ or $y = 3/2$. Since $3/2 > 1$, we have $y = 0$. \square

Proof of Theorem 5.1. First note that by adding the terms $(A'_{n,2})^3$ and $3A'_{n,2}(A'_{n,1})^2$ to the formula in Lemma 5.4, we obtain the estimate

$$A'_{n+1,1} \leq 54A_{n,1}^2 (A'_{n,1} + A'_{n,2}) - 9A_{n,1} (A'_{n,1} + A'_{n,2})^2 + (A'_{n,1} + A'_{n,2})^3.$$

Adding this to the formula in Lemma 5.5, we find that

$$\begin{aligned} |E_{n+1,\text{fix}}| &= A'_{n+1,1} + A'_{n,2} \leq 108A_{n,1}^2 (A'_{n,1} + A'_{n,2}) - 9A_{n,1} (A'_{n,1} + A'_{n,2})^2 + (A'_{n,1} + A'_{n,2})^3 \\ &= 3(6A_{n,1})^2 |E_{n,\text{fix}}| - \frac{3}{2}(6A_{n,1}) |E_{n,\text{fix}}|^2 + |E_{n,\text{fix}}|^3 \\ &= 3|E_n|^2 \cdot |E_{n,\text{fix}}| - \frac{3}{2}|E_n| \cdot |E_{n,\text{fix}}|^2 + |E_{n,\text{fix}}|^3, \end{aligned}$$

where we have used Lemma 5.2 to write $6A_{n,1} = |E_n|$. Dividing by $|E_{n+1}|$ and using Lemma 5.3, we find that

$$\frac{|E_{n+1,\text{fix}}|}{|E_{n+1}|} \leq \frac{|E_{n,\text{fix}}|}{|E_n|} - \frac{1}{2} \left(\frac{|E_{n,\text{fix}}|}{|E_n|} \right)^2 + \frac{1}{3} \left(\frac{|E_{n,\text{fix}}|}{|E_n|} \right)^3 = \phi \left(\frac{|E_{n,\text{fix}}|}{|E_n|} \right), \quad (5.5)$$

where ϕ is the polynomial from Lemma 5.6.

Set $x_n = \frac{|E_{n,\text{fix}}|}{|E_n|} \in [0, 1]$. To complete the proof, we must show that x_n tends to zero as $n \rightarrow \infty$. This follows immediately from (5.5) and Lemma 5.6, because

$$0 \leq x_{n+1} \leq \phi(x_n) \leq \phi^2(x_{n-1}) \leq \cdots \leq \phi^n(x_1) \leq \phi^n(1) \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad \square$$

6. ARITHMETIC APPLICATIONS

We now prove our applications on density of prime divisors in orbits and Newton's method. If K is a number field and \mathfrak{P} is a prime ideal of the ring of integers of K with residue field $k(\mathfrak{P})$, there is a surjective reduction map $K \rightarrow k(\mathfrak{P}) \cup \{\infty\}$. We write $x \equiv y \pmod{\mathfrak{P}}$ whenever $x, y \in K$ have the same reduction.

Proposition 6.1. *Let K be a number field and $x \in K$ an element such that (K, x) satisfies property (\dagger) of Section 3. Choose $y_0 \in K \setminus \{x\}$ and define a sequence $\{y_i\}_{i \geq 0} \subseteq K$ by $y_i = f^i(y_0)$. Then the set of prime ideals \mathfrak{P} of K such that*

$$y_i \equiv x \pmod{\mathfrak{P}} \quad \text{for some } i \geq 0$$

has natural density zero.

Proof. Note that for all $i \geq 0$, we have $y_i \neq x$. This inequality holds for $i = 0$ by hypothesis. Furthermore, if it failed for some $i > 0$, y_0 would be a K -rational solution to the equation $f^i(z) - x$, which is absurd since this polynomial is irreducible over K by Lemma 3.3.

For each $n \geq 1$, define S_n to be the set of prime ideals \mathfrak{P} of K such that

- x is not integral at \mathfrak{P} , or
- $y_i \equiv x \pmod{\mathfrak{P}}$ for some $0 \leq i \leq n-1$.

Then S_n is finite.

Let $n \geq 1$. Let $\mathfrak{P} \notin S_n$ be a prime ideal of the ring of integers of K such that $y_i \equiv x \pmod{\mathfrak{P}}$ for some $i \geq n$. Then $f^n(y_{i-n}) \equiv x \pmod{\mathfrak{P}}$, and therefore the polynomial $f^n(z) - x$ has a $k(\mathfrak{P})$ -rational root. Write $\delta(S)$ for the natural density of a set of prime

ideals S (if it exists). For each $n \geq 1$, the Chebotarev Density Theorem and the finiteness of S_n yield

$$\begin{aligned} & \delta(\{\mathfrak{P} \mid y_i \equiv x \pmod{\mathfrak{P}} \text{ for some } i \geq 0\}) \\ & \leq \delta(\{\mathfrak{P} \mid \mathfrak{P} \notin S_n, f^n(z) - x \text{ has a } k(\mathfrak{P})\text{-rational root}\}) \\ & = \frac{\left| \left\{ s \in \text{Gal}(K(f^{-n}(x))/K) \mid s \text{ fixes some root of } f^n(z) - x \right\} \right|}{|\text{Gal}(K(f^{-n}(x))/K)|} = \frac{|E_{n,\text{fix}}|}{|E_n|}. \end{aligned}$$

The final equality uses Theorem 3.7 to identify the Galois group of $f^n(z) - x$ over K with E_n . By Theorem 5.1, this last quantity tends to zero as $n \rightarrow \infty$. \square

Recall that the critical points for f are $0, 1, \infty$, and that they are all fixed by f .

Corollary 6.2. *Let K be a number field for which there exist unramified primes above 2 and above 3. Let $y_0 \in K \setminus \{0, 1, 3/2, -1/2\}$, and define a sequence $\{y_i\}_{i \geq 0} \subseteq K$ by $y_i = f^i(y_0)$. Then the set of prime ideals \mathfrak{P} of K such that*

$$y_i \equiv 0 \text{ or } 1 \pmod{\mathfrak{P}} \quad \text{for some } i \geq 0$$

has natural density zero. In particular, the set of prime divisors of the sequence $\{y_i\}_{i \geq 0}$ has natural density zero.

Proof. We begin by showing that the set of primes \mathfrak{P} such that $y_i \equiv 0 \pmod{\mathfrak{P}}$ for some $i \geq 0$ has natural density zero. Let S be the set of primes \mathfrak{P} of K for which

- \mathfrak{P} lies above 2, or
- $y_0 \equiv 0 \pmod{\mathfrak{P}}$.

Notice that since $y_0 \neq 0$, S is a finite set, and we may safely ignore primes in S for the remainder of the proof.

Suppose now that $\mathfrak{P} \notin S$ is such that $y_i \equiv 0 \pmod{\mathfrak{P}}$ for some $i \geq 1$. We may assume without loss that i is minimal with this property. Then

$$y_i = f(y_{i-1}) = -2y_{i-1}^3 + 3y_{i-1}^2 = -2y_{i-1}^2 \left(y_{i-1} - \frac{3}{2} \right),$$

which implies that $y_{i-1} \equiv \frac{3}{2} \pmod{\mathfrak{P}}$. We claim that $y_{i-1} \neq \frac{3}{2}$. This is true by hypothesis if $i = 1$. If it were to fail for some $i > 1$, then the polynomial $f^{i-1}(z) - 3/2$ would have y_0 as a K -rational root. But our hypothesis on K implies that $(K, 3/2)$ satisfies property (\dagger) , and so we have a contradiction to Lemma 3.3. It follows that

$$\delta(\{\mathfrak{P} \mid y_i \equiv 0 \pmod{\mathfrak{P}} \text{ for some } i \geq 0\}) = \delta\left(\left\{\mathfrak{P} \mid y_i \equiv \frac{3}{2} \pmod{\mathfrak{P}} \text{ for some } i \geq 1\right\}\right).$$

Since $(K, 3/2)$ satisfies property (\dagger) , the density on the right is zero by Proposition 6.1.

Now we show that the density of primes \mathfrak{P} such that $y_i \equiv 1 \pmod{\mathfrak{P}}$ for some $i \geq 0$ also has natural density zero. Define $w_i = 1 - y_i$ for $i \geq 0$. As $y_0 \notin \{1, -1/2\}$, we see that $w_0 \notin \{0, 3/2\}$. Moreover, because $1 - f(z) = f(1 - z)$, we find that

$$f(w_i) = f(1 - y_i) = 1 - f(y_i) = 1 - y_{i+1} = w_{i+1}.$$

Thus, we may apply the first part of the proof to the sequence $\{w_i\}_{i \geq 0}$ to deduce that

$$\delta(\{\mathfrak{P} \mid w_i \equiv 0 \pmod{\mathfrak{P}} \text{ for some } i \geq 0\}) = 0.$$

Since $w_i \equiv 0 \pmod{\mathfrak{P}} \Leftrightarrow y_i \equiv 1 \pmod{\mathfrak{P}}$, we are done. \square

We now prove a special case of the Faber-Voloch conjecture. Recall that the *Newton map* associated to a polynomial $g(z) \in K[z]$ is the rational function

$$N_g(z) = z - \frac{g(z)}{g'(z)}.$$

The simple roots of g are critical fixed points of N_g . Hence, for any completion K_v of K , the roots of g are super-attracting fixed points of the map N_g , viewed as a dynamical system acting on $\mathbb{P}^1(K_v)$.

Corollary 6.3. *Let K be a number field for which there exist unramified primes above 2 and above 3. Let $g(z) = z^3 - z$. Choose $y_0 \in K$ such that the Newton iteration sequence $y_i = N_g^i(y_0)$ does not encounter a root of g . Then the set of primes \mathfrak{P} of K for which the Newton sequence $\{y_i\}_{i \geq 0}$ converges in $K_{\mathfrak{P}}$ to a root of g has natural density zero.*

Proof. The Newton map for g is

$$N_g(z) = \frac{2z^2}{3z^2 - 1}.$$

Let $\eta(z) = 1/(1 - 2z)$. Then

$$\eta^{-1} \circ N_g \circ \eta(z) = -2z^3 + 3z^2 = f(z). \quad (6.1)$$

For each $i \geq 0$, define $w_i = \eta^{-1}(y_i)$. Then it is immediate from (6.1) that $w_{i+1} = f(w_i)$. Moreover, if we had $w_0 \in \{0, 1, 3/2, -1/2, \infty\}$, then the sequence $\{w_i\}_{i \geq 0}$ would encounter a fixed point of f , in which case $\{y_i\}_{i \geq 0}$ would encounter a fixed point of N_g and, hence, a root of g , contradicting our hypotheses. Thus, $w_0 \notin \{0, 1, 3/2, -1/2, \infty\}$. Corollary 6.2 therefore shows that the set of prime ideals \mathfrak{P} for which $w_i \equiv 0$ or $1 \pmod{\mathfrak{P}}$ has density zero.

On the other hand, the proof of the main theorem of Faber-Voloch [3] shows that for all but finitely many prime ideals \mathfrak{P} of K , the sequence $\{y_i\}_{i \geq 0}$ converges in $K_{\mathfrak{P}}$ to a root of g if and only if $g(y_i) \equiv 0 \pmod{\mathfrak{P}}$ for some $i \geq 0$. Factoring g , this condition is equivalent to saying that $y_i \equiv 0, \pm 1 \pmod{\mathfrak{P}}$ for some $i \geq 0$, which in turn is equivalent to saying that $w_i \equiv 0, 1$, or $\infty \pmod{\mathfrak{P}}$. (Here, $w \equiv \infty \pmod{\mathfrak{P}}$ means w is not integral at \mathfrak{P} .) Clearly, the set of primes \mathfrak{P} for which $w_i \equiv \infty \pmod{\mathfrak{P}}$ is zero, since f is a polynomial, and so the proof is complete. \square

Acknowledgments: This project began at a workshop on “The Galois theory of orbits in arithmetic dynamics” at the American Institute of Mathematics in May 2016. The authors would like to thank AIM for its generous support and hospitality, Rafe Jones for his early encouragement to pursue this line of thought, and Clay Petsche for several early discussions. The first author gratefully acknowledges the support of NSF grant DMS-1501766. The third author gratefully acknowledges the support of NSF grant DMS-1415294.

REFERENCES

- [1] Robert L. Benedetto, Dragos Ghioca, Benjamin Hutz, Pär Kurlberg, Thomas Scanlon, and Thomas J. Tucker. Periods of rational maps modulo primes. *Mathematische Annalen*, pages 1–24, 2012.
- [2] Michael R. Bush, Wade Hindes, and Nicole R.Looper. Galois groups of iterates of some unicritical polynomials. [arXiv:1608.03328v1](https://arxiv.org/abs/1608.03328v1) [math.NT], preprint, 2016.
- [3] Xander Faber and José Felipe Voloch. On the number of places of convergence for Newton’s method over number fields. *J. Théor. Nombres Bordeaux*, 23:387–401, 2011.

- [4] Richard Gottesman and Kwokfung Tang. Quadratic recurrences with a positive density of prime divisors. *Int. J. Number Theory*, 6(5):1027–1045, 2010.
- [5] Rafe Jones. The density of prime divisors in the arithmetic dynamics of quadratic polynomials. *J. Lond. Math. Soc. (2)*, 78(2):523–544, 2008.
- [6] Rafe Jones. Galois representations from pre-image trees: an arboreal survey. *Publ. Math. Besançon*, pages 107–136, 2013.
- [7] Rafe Jones and Michelle Manes. Galois theory of quadratic rational functions. *Comment. Math. Helv.*, 89(1):173–213, 2014.
- [8] J. Juul. Iterates of generic polynomials and generic rational functions. [arXiv:1410.3814v4 \[math.NT\]](#), preprint, 2016.
- [9] Nicole R.Looper. Dynamical Galois groups of trinomials and Odoni’s conjecture. [arXiv:1609.03398v1 \[math.NT\]](#), preprint, 2016.
- [10] R. W. K. Odoni. The Galois theory of iterates and composites of polynomials. *Proc. Lond. Math. Soc. (3)*, 51(3):385–414, 1985.
- [11] R. W. K. Odoni. On the prime divisors of the sequence $w_{n+1} = 1 + w_1 \cdots w_n$. *J. Lond. Math. Soc. (2)*, 32(1):1–11, 1985.
- [12] Richard Pink. Profinite iterated monodromy groups arising from quadratic polynomials. [arXiv:1307.5678 \[math.GR\]](#), preprint, 2013.

(Benedetto, Juul) AMHERST COLLEGE, AMHERST, MA
E-mail address, Benedetto: rlbenedetto@amherst.edu

(Faber) CENTER FOR COMPUTING SCIENCES, INSTITUTE FOR DEFENSE ANALYSES, BOWIE, MD
E-mail address, Faber: awfaber@super.org

(Hutz) SAINT LOUIS UNIVERSITY, SAINT LOUIS, MO
E-mail address, Hutz: hutzba@slu.edu

(Juul) AMHERST COLLEGE, AMHERST, MA
E-mail address, Juul: jamie.1.rahr@gmail.com

(Yasufuku) COLLEGE OF SCIENCE AND TECHNOLOGY, NIHON UNIVERSITY, TOKYO, JAPAN
E-mail address, Yasufuku: yasufuku@math.cst.nihon-u.ac.jp